

Assistant Professor of Orthopaedics Division of Sports Medicine Tel: (646)501-7122

Iliotibial Band Friction Syndrome Stretching Exercises


Name:	Date:
Diagnosis:	Date of Surgery:
■ TABLE HANG STRETCH • Lie on your side with the uninvolved thigh on the table ar While keeping your involved leg straight, extend the leg a gravity pull the involved leg to the floor. Hold second	at the hip so that it hangs over the table edge. Let
 LEAN AWAY OR WALL STRETCH Stand with both legs completely straight. Place the involved as far as possible behind the uninvolved leg. Lean or push Hold seconds. Repeat times. 	
 LEAN AWAY WITH TRUNK FLEXION STRETCH Stand with both legs completely straight. Place the involved knee is directly behind the uninvolved knee. Leg forward as far as possible. Hold seconds. Repeat 	an away from the involved leg as you bend

Guillem Gonzalez-Lomas, MD

Hospital for Joint Diseases

NYU LANGONE MEDICAL CENTER

Assistant Professor of Orthopaedics Division of Sports Medicine Tel: (646)501-7122



• Lie on the involved leg with your knees and hips in a straight line with the trunk. Press up with your arm directly under your shoulder, supporting your trunk with your extended arm. Hold seconds. Repea times.
• Sit with your involved leg bent and crossed over the top of your uninvolved leg. Rotate your trunk toward your involved leg. Place your elbow on the outside of your involved leg. Gently push your involved leg away as far as possible using your elbow. Hold seconds. Repeat times.
Comments:
Frequency: times per week
Signature: Date: